

Corrigendum

Morphological distinction between vasodilator and secretomotor neurons in the pterygopalatine ganglion of the cat [Neurosci. Lett. 288 (2000) 219–222][☆]

Satoshi Kuchiiwa^{a,*}, Shi-Bin Cheng^a, Toshiko Kuchiiwa^b

^a*Department of Anatomy, Faculty of Medicine, Kagoshima University, Sakuragaoka 8-35-1, Kagoshima 890-8520, Japan*

^b*Department of Life Science, Kagoshima Immaculate Heart College, Toso 4-22-1, Kagoshima 890-8525, Japan*

Received 7 March 2000; received in revised form 18 May 2000; accepted 1 June 2000

In the original article, the last line of the abstract read “Thus, it was assumed that small and large neurons labeled in the ganglion represented, respectively, secretomotor and vasomotor neurons.”

The correct form is “...represented, respectively, vasomotor and secretomotor neurons.”

Also in the original article, reference [1] was incorrect - the correct version is printed below.

- [1] Cheng, S.-B., Kuchiiwa, S., Kuchiiwa, T. and Nakagawa, S., Three novel neural pathways to the lacrimal glands of the cat: an investigation with cholera toxin b subunit as a retrograde tracer, *Brain Res.*, 873 (2000) 160–164.

[☆] PII of original article: S0304-3940(00)01236-2.

* Corresponding author. Tel./fax: +81-99-275-5214.

E-mail address: kuchiiwa@med1.kufm.kagoshima-u.ac.jp (S. Kuchiiwa).